

§ 522.275

use by or on the order of a licensed veterinarian.

(2) *Horses*—(i) *Amount*. 0.05 milligram of butorphanol base activity per pound of body weight (0.1 milligram/kilogram) using 10 milligrams per milliliter solution.

(ii) *Indications for use*. For the relief of pain associated with colic and postpartum pain in adult horses and yearlings.

(iii) *Limitations*. For intravenous use in horses only. Dose may be repeated within 3 to 4 hours. Treatment should not exceed 48 hours. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(3) *Cats*—(i) *Amount*. 0.2 milligram of butorphanol base activity per pound of body weight (0.4 milligram/kilogram), using 2 milligrams per milliliter solution.

(ii) *Indications for use*. For the relief of pain in cats caused by major or minor trauma, or pain associated with surgical procedures.

(iii) *Limitations*. For subcutaneous injection in cats only. Dose may be repeated up to 4 times per day. Do not treat for more than 2 days. Safety for use in pregnant female cats, breeding male cats or kittens less than 4 months of age has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 29276, May 2, 1980, as amended at 50 FR 24508, June 11, 1985; 53 FR 27851, July 25, 1988; 59 FR 41665, Aug. 15, 1994; 63 FR 66432, Dec. 2, 1998; 68 FR 22594, Apr. 29, 2003]

§ 522.275 N-Butylscopolammonium bromide.

(a) *Specifications*. Each milliliter of solution contains 20 milligrams (mg) N-butylscopolammonium bromide.

(b) *Sponsor*. See No. 000010 in § 510.600(c) of this chapter.

(c) *Conditions of use in horses*—(1) *Amount*. 0.3 mg per kilogram of body weight (0.14 mg per pound) slowly intravenously.

(2) *Indications for use*. For the control of abdominal pain (colic) associated with spasmodic colic, flatulent colic, and simple impactions.

21 CFR Ch. I (4–1–05 Edition)

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[69 FR 35512, June 25, 2004]

§ 522.311 Carfentanil citrate injection.

(a) *Specifications*. Each milliliter of sterile aqueous solution contains 3 milligrams of carfentanil citrate.

(b) *Sponsor*. See No. 053923 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 5 to 20 micrograms per kilogram (.005 to .020 milligram per kilogram) of body weight.

(2) *Indications for use*. For immobilizing free ranging and confined members of the family Cervidae (deer, elk, and moose).

(3) *Limitations*. Inject into large muscle of neck, shoulder, back, or hindquarter. Avoid intrathoracic, intra-abdominal, or subcutaneous injection. To reverse effect, use 7 milligrams of diprenorphine for each milligram of carfentanil citrate, given intravenously or one-half intravenously and one-half intramuscularly or subcutaneously. Do not use in domestic animals intended for food. Do not use 30 days before or during hunting season. Do not use in animals that display clinical signs of severe cardiovascular or respiratory disease. Available data are inadequate to recommend use in pregnant animals. Avoid use during breeding season. Federal law restricts this drug to use by or on the order of a licensed veterinarian. The licensed veterinarian shall be a veterinarian engaged in zoo and exotic animal practice, wildlife management programs, or research.

[53 FR 40057, Oct. 13, 1988]

§ 522.312 Carprofen.

(a) *Specifications*. Each milliliter of solution contains 50 milligrams (mg) carprofen.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use in dogs*—(1) *Amount*. 2 mg/lb (4.4 mg/kg) body weight once daily or 1 mg/lb (2.2 mg/kg) twice daily, by subcutaneous injection. For the control of postoperative pain,